



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

glands, would seem naturally to suggest some theoretical views which may prove of use as a guide for future investigations.

As the two styles in Apocynum and Asclepias are more or less distinct both above and below, and each of them belongs to a simple pistil, it is not easy to account for a five angled stigma with five pairs of glands. But if, as De Candolle supposes, the diaphragm in Apocynum is an extension of the filaments which has become attached to and surrounds the styles, its fivefold character is readily accounted for in Apocynum, and, in view of the close relationship of the Orders and the similarity of their fertilization, it would, perhaps, not be unreasonable to extend his supposition to what is called the peltate stigma of Asclepias.

It has been shown that the glands in Apocynum arise in pairs from the middle of the lobe of the diaphragm, where it joins the filament. It is possible, therefore, that these glands are likewise a modification of some organ of the stamen. Each half stretches away from the point of origin of the pair towards the adjacent cell of its own anther, but when mature, and ready to be carried away with the pollen, it lies so near to the half of the next pair on the right or left, that a new pair seems to be made up from and between adjacent stamens. Now Brown has shown that the dark glands in Asclepias consist at first of two distinct parts, and the halves are really related to different anthers. The parallelism of the cases may justify the hypothesis that these glands of Asclepias are members of the stamen, if that should prove to be true of Apocynum. Perhaps a clue to the glands is the pollen missing in the lower part of the anther of Apocynum.

On this theory, which is only an extension of what is already recognized, the so-called stigma and the glands of Asclepias are merely a modification of the stamens, arising from the adhesion of the latter to the style. The peculiarity of such adhesion in these two Orders is that it takes place in the upper instead of the lower part of the organs.

W. H. L.

4. *Lygodium palmatum*, Swartz.—I have found the beautiful *Climbing Fern* growing in the drier parts of a sphagnum bog, some two miles westerly from Mount Pleasant, which is a little southerly from Matawan, Monmouth Co., N. J. The specimens were very fine and full of fruit.

S. LOCKWOOD.

5. *Aquilegia*.—In the *Flowering Plants of Great Britain*, by Anne Pratt, we find the following: "The French term this plant *L'Ancolie*; and it is *Der Akeley* of the Germans. The Italians call it *Acquilezia*." In the Penny Cyclopædia we find: "*Aquilegia*, literally the *Watergatherer*, because the leaves collect water in their hollow." A friend informs us that he has noticed this peculiarity.

6. *Exchanges*.—J. W. Congdon of East Greenwich, R. I., desires to make arrangements for exchange of collections to be made the following season. He will collect all the less common plants of Rhode Island, and of Northern Vermont, including the Willoughby plants. He has quite a number of them in hand now.

He particularly desires to secure New Jersey, Southern and Western plants.

7. New Publications.—*The American Journal* for Dec. contains an instructive notice by Dr. Gray of Decaisne's *Monograph of the Genus Pyrus*.—*The American Naturalist* for Dec., appetizing articles on *The Vegetation of the Lower Wabash Valley*, and the *Alpine Flora of Colorado*; for Jan., Mr. Bailey on *Our Poisonous Plants*, and Mr. Biscoe on *The Study of Lichens*. We recommend all our readers to subscribe to this most deserving Journal. We notice an oversight in not crediting us with Mr. Merriam's article on *Trees and Rain*.—*The Popular Science Monthly* has an article on *Cultivating Wild Flowers*, by Rev. S. Lockwood.—*Nature*, Oct. 10 and 17, has some very interesting observations on the *Fertilization of a few common Papilionaceous Flowers*.

ORCHIDACEÆ.

ORCHIS, L.—*O. spectabilis*, L.; not rare; High Bridge; Hoboken; New Durham, etc.

HABENARIA, Willd., R. Br.—*H. tridentata*, Hook.; not uncommon; New Durham and Secaucus swamps; Closter, *Austin*; Big Swamp, Chatham, N. J.; Staten Island, *W. H. L.*—*H. virescens*, Spreng.; not uncommon; Closter, Secaucus, Chatham, N. J.; Yonkers, *Pooley*; Glen Cove, *Coles*.—*H. viridis*, R. Br., var. *bracteata*, Reichenbach; Closter, very rare, *Austin*.—*H. Hookeri*, Torr.; Westchester Co., *Le Roy*; abundant in Orange Co., *Austin*.—*H. orbiculata*, Torr.; Closter, very rare, *Austin*.—*H. cristata*, R. Br.; New Durham Swamp, *Torr. Cat.*, but not reported of late years.—*H. ciliaris*, R. Br. In a swamp in Love-lane, and in Hoboken meadows, *Torr. Cat.*; Tappan and Tennaflly, *Austin*; Staten Island and Redbank, N. J., *W. H. L.*; Keyport, N. J., *R. W. Brown*; Glen Cove, *Coles*.—*H. blephariglottis*, Hook.; Keyport, *Brown*; Closter, *Austin*; Secaucus, *W. H. L.*—*H. lacera*, R. Br.; common in all our districts; N. Y.—*H. psycodes*, Gray; common; Closter, *Austin*; Glen Cove, *Coles*; Chatham, N. J.; *W. H. L.*—*H. fimbriata*, R. Br.; "Meadows about Elizabethtown, N. J.," *Eddy in Torr. Cat.*

GOODYERA, R. Br.—*G. pubescens*, R. Br.; common; N. Y.

SPIRANTHES, Richard.—*S. latifolia*, Torr.; reported from Glen Cove, but needs confirmation.—*S. cernua*, Richard; common.—*S. graminea*, Lindl., Var. *Walteri*; common; Staten Island and Summit, N. J., *W. H. L.*; Closter and Morris Co., N. J., *Austin*; Train's Meadow, Newtown, L. I., *Ruger*.—*S. gracilis*, Bigelow; the most common form in all our districts; N. Y.—*S. simplex*, Gray; Tottenville, Staten Island, rare, *W. H. L.*; Keyport, quite common, *Brown*; Closter, rare, *Austin*.

LISTERA, R. Br.—*L. cordata*, R. Br.; probably grows in our region, though no specimen properly authenticated is at hand.—*L. convallarioides*, Hook.; "In the cedar swamp at New Durham," *Torr. Cat.*

ARETHUSA, Gronov.—*A. bulbosa*, L.; Closter, rare, *Austin*; New Lots, *Merriam*; Glen Cove, *Coles*; New Durham Swamp, formerly,